

Home preparation of juices, wines & cider

PUBLICATION 1406 1975
REVISED



630.4
C212
P 1406
1975
(1976 print)
c. 3



Agriculture
Canada

Copies of this publication may be obtained from
INFORMATION DIVISION
CANADA DEPARTMENT OF AGRICULTURE
OTTAWA
K1A 0C7

© MINISTER OF SUPPLY AND SERVICES CANADA 1976

Printed 1970

Reprinted 1970, 1972, 1974, 1976

Revised 1975

20M-38390-8:76

Cat. No.: A73-1406


ISBN 0-662-00151-6

Librairie Beauchemin Limitée
Contract No. 08KT-01A05-6-38390

CANADA DEPARTMENT OF AGRICULTURE PUBLICATION 1406 REVISED 1975

Home preparation of juices, wines, and cider

D.R. MacGregor, J.A. Ruck, and J.F. Bowen Revised by J.F. Bowen
Research Station, Summerland, British Columbia



CONTENTS

Juices

<i>Apple juice</i>	3
<i>Juices other than apple</i>	6
<i>Tomato juice</i>	6
<i>Grape juice</i>	7
<i>Apricot, prune, and peach juices</i>	8
<i>Black currant juice</i>	9
<i>Miscellaneous juices</i>	11
<i>Freezing juices</i>	11

Wines

<i>Units of measurement</i>	12
<i>Grape wine</i>	12
<i>Equipment</i>	12
<i>Preparing the pulp or juice</i>	12
<i>Yeasts and fermentation</i>	15
<i>Addition of sugar</i>	16
<i>Racking and fining the wine</i>	18
<i>Aging</i>	19
<i>Fruit wines</i>	19
<i>Apricot, peach, and prune wines</i>	20
<i>Cherry wine</i>	20
<i>Loganberry and raspberry wines</i>	20
<i>Black currant wine</i>	21
<i>Dandelion wine</i>	21
<i>Parsnip wine</i>	22
<i>Rhubarb wine</i>	22

Cider

<i>Containers for fermentation</i>	23
<i>Preparing the juice</i>	23
<i>Fermentation</i>	24
<i>Pasteurization</i>	24
<i>Storage</i>	24
<i>Still sweet cider</i>	25
<i>Things to Remember</i>	25
<i>Glossary of Terms</i>	25



More and more people are becoming interested in preparing various fruit juice products at home. Improvements in equipment and techniques have made the methods for preparing juices, wines, and cider both simple and practical.

juices

In making small quantities of juice, be sure to handle the fruit carefully and carry out the various steps efficiently. Limit the size of each batch to a quantity that can be handled quickly from the preparation of the fruit to the bottling and cooling of the juice. To prevent loss in nutritive value, discoloration, and even spoilage of the juice, do not heat the juice any longer than necessary during preparation, and fill the containers as quickly as possible. Be sure the containers are thoroughly clean before you start.

APPLE JUICE

Use only mature, sound apples for making juice. All commonly grown varieties with the exception of Delicious and the very early summer apples give a juice of good quality. Juices from the Delicious variety are too low in acid to be used alone, therefore blend them with more acid juices in amounts not exceeding 20 to 25%.

A pulpy type of apple juice can be made by using almost any home juice machine, but if a clearer juice is desired a hammer mill and press must be used (Figures 1 to 5). This equipment is easy to use, and for economy a number of families can share the cost. The drawing and general description of the juice press

(Figure 3) have been adapted from Circular 194 of the New York Agricultural Station, Geneva, N.Y. Contact with iron or copper may cause discoloration, off-flavor, and cloudiness, therefore, all equipment coming in contact with the juice should be made of stainless steel, glass, or plastic (Figure 6). Some plastic paints are suitable for coating iron equipment.

1. Mill and press the fruit, and strain the juice through clean cheesecloth. Juice extracted in this way is cloudy, or "opalescent," and may be used without further clarification. However, if a clear juice is desired add the enzyme pectinase at the rate of $\frac{1}{4}$ teaspoon/gal ($1 \text{ cm}^3/4 \text{ litres}$). Allow the treated juice to stand overnight and then siphon the clear liquid free from the sediment. Pectinase is available under various trade names, such as Pectozyme and Clarase, from stores carrying home wine makers' supplies and some food specialty and drug stores. It should not be confused with pectin, the material used in making jellies.

Juices may be fortified with vitamin C by adding ascorbic acid at the rate of $\frac{1}{2}$ teaspoon/gallon ($2 \text{ cm}^3/4 \text{ litres}$) just before the containers are filled. Dissolve it in a small amount of juice or water, add this to the main batch, and stir well. This treatment helps to preserve the color and gives a vitamin C content equivalent to that of most citrus juices. The juice may be stored without further treatment in the home deep freezer. Cardboard milk cartons make useful containers, if they are cleaned well.

2. Heat the clarified or unclarified juice to 200°F (93°C) in an acid-resistant container.
3. To prepare plain tin cans, wash them thoroughly in hot water, rinse with boiling water, and invert to drain. Pour the hot juice into the cans, seal the cans, and invert them for 2 to 3 min. Cool the juice by putting the cans into cold water immediately. Fast cooling gives a better-flavored juice.
4. If glass sealers are used, wash them and sterilize by heating in an oven to $190 - 200^\circ\text{F}$ ($88 - 93^\circ\text{C}$), or use an automatic dishwasher set on the sanitizing cycle. Boil or scald the lids. Fill the containers completely, seal or cap them and stand them in water at 110°F (43°C) until cool. Place crown-capped bottles on their sides for a few minutes before




Figure 1. Apple mill.

cooling to sterilize the caps.

5. Store the filled containers upright in a cool place.
Juices prepared in this way develop some sediment during

storage, but when they are opened and the liquid is poured off carefully, most of the sediment remains in the bottom of the container. This sediment is completely wholesome and affects only the appearance of the juice.

JUICES OTHER THAN APPLE

Juices from fruits other than apple usually contain some pulp, and the equipment for making them (Figure 7) is much simpler. Several types of juice extractors do a good job. The best one operates on the principle of a tapered screw that presses the juice through a surrounding tapered screen and extrudes the waste material from the end. Another one extracts the juice by steam, but yields a diluted juice with a cooked flavor. In another type the fruit contacts a rapidly spinning toothed disc and the juice is expelled by centrifugal force. Unfortunately, this type tends to incorporate air into the juice. The cone-shaped food press is the cheapest, but a great deal of work is involved in making a relatively small amount of juice.

Expensive equipment could be purchased cooperatively by several families or by groups such as a Women's Institute Chapter or Wine Guild.

Tomato Juice

1. Stem fully ripe tomatoes, and trim them to remove all moldy material.
2. Pulp the tomatoes in a preserving kettle and bring the pulp to a boil as quickly as possible.
3. Pass the hot pulp through a juice extractor or sieve to remove the skins and seeds.
4. Sterilize sealers or bottles as described for apple juice.
5. Add salt at 1 tablespoon/gal of extracted juice (12 cm³/4 litres) and reheat the juice to the boiling point.
6. Quickly pour the juice into sealers, bottles, or cans, and seal the containers; do not leave any headspace. Excessive air in the headspace causes serious loss of vitamin C during storage. The juice must be at a temperature of at least 200°F (93°C) when the containers are being filled.

7. Invert cans for 3 to 5 min to sterilize the lids, and then cool them in cold water. Place bottles on their sides to sterilize the caps, and air-cool them, or water-cool them as in step 4 for apple juice. Sealers must be cooled in the upright position in order for a proper seal to be made; because the lids will have been sterilized beforehand, inverting the jars is unnecessary.

Grape Juice

For the best flavor use fully ripened grapes, free from mold. Concord type varieties give a fruity, highly flavored juice. Varieties such as New York Muscat give a strongly muscat-flavored juice, which may be diluted to taste with juice from neutral-flavored grapes, or with apple juice.

1. Remove the stems and crush the grapes enough to release the juice but not to break the seeds.
2. Heat the pulped fruit to 145-150°F (63-66°C) to extract color and other soluble materials. Avoid prolonged heating because it extracts excessive tannin from the seeds and skins, which may cause bitterness or off-flavors.
3. After heating, press the pulp in a juice press (Figures 3,4, and 5) or pass it through a juice extractor (Figure 7) to remove the skins and seeds.
4. When necessary, strain the juice through cheesecloth to remove the large particles of pulp. If storage at 28-30°F (-2 to -1°C) is available, hold the juice at this temperature for 24-48 hr to allow sediment to settle out. Pectinase (see page 4) may be added at this point to speed clarification.
5. Siphon the clear juice free from the sediment and heat to 180-185°F (82-85°C) for 3 min. Dispense into clean, airtight bottles. Cap them immediately and cool as in step 4 for apple juice. After prolonged storage some tartrate (cream of tartar) may settle out. When the bottles are opened, pour the juice carefully so that the sediment is not disturbed.

Freshly pressed unheated juice may be stored in milk car-

tons in the deep freezer. For this purpose, the grapes may be pressed cold. Add one Campden tablet (see page 13), pectinase, and ascorbic acid at $\frac{1}{4}$ teaspoon/gal ($1 \text{ cm}^3/4 \text{ litres}$) to preserve color and allow the juice to settle for several days to one week at $28\text{-}30^\circ\text{F}$ (-2 to -1°C). The clear juice may then be drawn off and dispensed into the containers and frozen.

Apricot, Prune, and Peach Juices

These juices are made in much the same way as tomato juice. Use fully ripe fruit to obtain the best color and flavor.

1. Prepare the apricots and prunes by halving, pitting, and pulping. Remove peach skins by using a hot-water blanch




Figure 2. Apple mill with hopper open. A, 1/2 hp heavy-duty motor, 1750 rpm, 5/8-in. shaft. B, 5/8-in. flexible coupling. C, Steel blades $\frac{3}{16} \times 1 \frac{1}{2} \times 7 \frac{3}{4}$ in. straight ends. D, 3/4-in. shaft threaded for locking nuts, G. The thread on one end is reversed so that the locking nuts tighten when the blades rotate. Both ends of the shaft are turned down to 5/8 in. Spacers between blades are $\frac{7}{16}$ in. long and can be cut from 3/4-in. pipe. E, 5/8-

in. self-aligning permanently lubricated bearings, which lock to the shaft. F, Half-round 8-in. diam 12-gauge steel screen perforated with 1/2-in. holes. The holes are drilled on the points of equilateral triangles that have 7/8-in. sides. G, Locking nuts. H, Hopper. Offset 22° from the vertical. Throat is 4 x 8 in. 12-gauge. Section around the screen is 8 x 8 in. Discharge below the mill is 7 x 7 in. Height of discharge to floor is 12 in.

as in canning, and then halve, pit, and pulp the peaches.

2. Place the prepared fruit in a suitable preserving kettle, add water at 2 cups to each 5 lb of fruit (200 cm³/kg) and stirring constantly, bring the mixture rapidly to a boil.
3. Pass the hot pulp through a juice extractor to remove skins and fibres.
4. To make the extracted juice into a full-bodied drink, add sugar at 1 lb to each 5 gal of juice (0.4 kg/20 litres). To make a lighter product, dilute the extracted juice with an equal amount of a syrup prepared by dissolving 1 lb of sugar in ½ gal of water (0.2 kg/litre).
5. Reheat the juice to boiling and pour it into sterilized containers. Sterilize the containers and cool the juice as in step 4 for apple juice.

For peach juice with a distinctive flavor, to each 4 cups (litre) of undiluted juice add 1–2 tablespoons (16–31 cm³) of frozen orange concentrate and 1–2 teaspoons (5–10 ml) of lemon juice or 1 tablespoon (16 ml) of lime juice, or 2–3 tablespoons (31–47 cm³) of frozen grapefruit concentrate, according to taste.

Apricot juice blended with apple juice makes a pleasantly flavored drink.

Black Currant Juice

Black currants, an excellent source of vitamin C, make a tasty fruit juice.

1. In a suitable kettle, add 1 cup of water to each pound of black currants (0.5 litres/kg) and, stirring constantly, bring the mixture to a boil.
2. Carry out the extraction, reheating, filling, capping, and cooling as outlined from step 3 for tomato juice.




Figure 3. Juice press. A, Hydraulic jack, 1 1/2 tons capacity, 8-in. lift. B, Jack support, a 14 x 14-in. piece of 3/4-in. fir plywood or hardwood with 6 slats 1/4 x 1 in. nailed on one side. C, Racks. Five racks 14 x 14 in. are made from 1/4 x 1-in. slats of hardwood nailed to 1/4 x 1 1/2-in. slats at the edges. The center slat is 18 in. long and serves as a guide between the uprights, I. It is 2 in. wide and has a center slot. The nails are stainless steel. D, Press cloths of medium factory cotton are about 36 in. square. E, Press base is a 17 x 17-in. piece of 3/4-in. plywood fastened to supports. G, A 14 x 14-in. piece of 1/2-in. plywood is centered on the larger piece to channel the juice to the outlet, F. Four slats 1/4 x 1 1/2 in. on edge form the sides around the base. F, Juice outlet made of 3/4-in., acid-resistant or plastic pipe. G, Supports for press base. H, Press top is of hardwood or fir plywood with a metal plate at the point of contact of the jack. I, Uprights are reinforced with 1/4 x 4-in. metal strapping across the press top and 20 in. down two sides. All uprights and crosspieces are made of 4 x 4-in. fir. J, Cheese form is 2 x 14 x 14 in. and is made from 3/4-in. hardwood. All wooden parts of the press that come in direct contact with the juice are coated with hot paraffin.

Boil new cotton cloths in water for 5 min. and rinse. Dampen them before use. After each use, wash the cloths thoroughly by changing the water often and dry them well before you store them. To use the press, place a rack on the press base. Put the cheese form on the rack and cover both with a press cloth. Pour the product to be pressed on the cloth, as in Figure 5, to make a cheese 2 in. thick. Fold the cloth neatly into

the center of the cheese and remove the cheese form. Place another rack above this cheese and repeat the procedure until there is only space for the jack support, B, and the jack. Center the jack on the jack support and apply pressure evenly.

3. Before reheating and filling, dissolve 7 lb of sugar in each gallon of juice (3.6 kg / 5 litres).
4. Before drinking the juice, dilute it with an equal quantity of water. A delicious drink can be made by mixing one can of black currant juice with three cans of apple juice.
5. Store black currant juice at a temperature below 77°F (25°C) in the dark, so that it will retain its attractive color and delicate flavor.

Miscellaneous Juices

Appetizing juices may be made from many of the less common wild and cultivated fruits by following the general principles outlined here. Because of the danger of spoilage and loss of flavor, low-acid vegetable juices such as carrot and cabbage should not be preserved by heat, but they may be frozen. However, since these vegetables store well, it is much better to make fresh juice when it is wanted.

FREEZING JUICES

Any of the above juices may be frozen and stored in a home freezer. Cool the juice quickly after it has been extracted from the heated pulp. Pour the juice into suitable freezer containers or containers with plastic liners. Paper milk cartons can be washed and used for this purpose. Fill the containers only three-quarters full to allow for expansion during freezing.

wines

Wine is usually made by fermenting the pulp or juice of grapes, but delicious wine can also be made from all common fruits and even some vegetables.

UNITS OF MEASUREMENT

The amateur wine maker is often confronted with some metric units of measurement. A table of conversion factors for the metric system is included inside of the back cover of this publication.

GRAPE WINE

Equipment

Use only glass, plastic, or earthenware utensils for fermenting the pulp or juice. Contact with iron or copper causes the wine to darken, and zinc-coated (galvanized) containers may produce unpleasant flavors. Avoid enamelware containers, particularly if the coating is chipped. Stoneware crocks are the most convenient for fermenting pulped fruit, and glass or plastic carboys are the best for fermenting juice. The narrow neck of the carboy permits the use of a fermentation-lock stopper, which does not allow air or fruit flies to enter the carboy.

Clean all utensils thoroughly and rinse them with boiling water before use. Wooden barrels, if used for storing wine, should be free from the musty or vinegary odors caused by molds and bacteria. Sterilize them by soaking for 24 hr with a solution prepared by dissolving 8-10 Campden tablets in 5 gal (23 litres) of water (see following section). After this treatment rinse the barrels thoroughly with water.

When storing barrels empty, keep them dampened and add a little of the above solution from time to time.

Preparing the Pulp or Juice

Grapes should be picked when they are fully ripe and the sugar content and flavor are at their peak. Remove the stems and crush the grapes to release the juice. Avoid crushing the seeds because this may cause bitterness in the finished wine. A wooden plunger may be used to crush large quantities of grapes.

Red wines are made by fermenting the skin, pulp, and juice together; the skins impart the red color.

For white wines, the juice is preferably extracted before fermentation, but a satisfactory white wine can be made from




Figure 4. Juice press showing the various parts. A, Jack. B, Bottom of jack support. C, Rack. D, Cheese form.

white grapes fermented with the skins left on. However, the juice should be separated from the pulp after no more than 24 hr fermentation for best results.

When all the fruit has been thoroughly crushed or the juice extracted, and measured, add 2 Campden tablets, $\frac{1}{4}$ teaspoon (1.2 cm^3) of ascorbic acid, and pectinase according to the directions on the package, to each gallon (4.5 litres) of juice or 10 lb (4.5 kg) of pulped fruit. Stir the mixture and allow it to stand at room temperature for 24 hr.



Figure 5. Filling the cheese.

Campden tablets are composed of potassium metabisulphite, which when added to the juice releases sulphur dioxide. This sterilizes the juice by eliminating wild yeasts, molds, and bacteria, which if present may cause an undesirable vinegary or musty flavor. It also helps to preserve the color and flavor of the wine. Campden tablets are sold by stores that specialize in wine producers' supplies. They are much more convenient to use than the powdered metabisulphite. However, if the powdered form is used, a scant $\frac{1}{4}$ teaspoon (1.2 cm^3) and no more is the equivalent of 2 tablets.

Ascorbic acid in conjunction with metabisulphite prevents oxidation and undesirable color and flavor changes in the wine.

Yeasts and Fermentation

Pure cultures of wine yeasts in the form of packaged dry yeast or tubed agar cultures may be bought at stores that specialize in wine producers' supplies. They are selected yeast strains, capable of growing in the presence of sulfur dioxide, able to produce high concentrations of alcohol, and give a clean-flavored wine. Their use minimizes the risk of having the wine spoiled by the formation of vinegar. Bread yeasts are not recommended as a wine culture because the amount of alcohol they produce is insufficient to preserve the wine. Bread yeasts may be used as a last resort if wine yeasts are not available.

For making 5 to 10 gal (23–45 litres) of wine, sprinkle one package of dry yeast over the surface of the must (crushed grapes or juice), and when islands of bubbles appear, stir the mixture and allow it to ferment at 65–75°F (18–24°C) in a dark place.

To use a tubed agar culture, first fill the tube containing the culture with juice that has been sterilized by boiling and cooled to about 100°F (38°C). As soon as froth or bubbles appear, which means that fermentation is starting, pour the liquid into a gallon (4.5 litres) of sulfited juice.

When making more than 10 gal (45 litres) of wine in one batch, it is best to prepare a starter. To do this, make a small batch of wine from either the packaged dry yeast or the agar culture. After about 24 hr, when this small batch is fermenting strongly, add it to the main lot of must. Use the starter in the proportion of not less than 1 gal to 25 gal of must (4 litres to 100 litres).

Pressed juice for white wines may be fermented to completion in a large glass bottle or carboy with a fermentation lock in place to exclude fruit flies and to allow gas to escape.

The fermentation of pulped fruit may be carried out in a crock covered with cheesecloth or plastic film. A heavy cap of skins will rise to the surface and must be stirred in at frequent intervals. After 3-4 days (24 hr in the case of white wines) siphon off the free run juice into a clean carboy for further fermentation. If the juice remains on the pulp too long, it will extract excessive flavor and bitterness from the seeds and skins. Apply only gentle pressure to the drained pulp to extract the wine remaining in it.

Addition of Sugar

The amount of alcohol and sweetness in the finished wine depends on the amount of sugar in the juice. If the grapes are low in sugar, extra sugar may be added before fermentation.

The total sugar content, including the natural sugar present in the grapes, should not be more than 20% for good dry table wine. If more than this amount is present, the wine will have a



Figure 6. Small equipment and supplies. A, Five-gallon glass or plastic carboy. B, Four feet of 1/2-in. plastic or rubber tubing. C, Beer bottles for cider or juice. D, Capper. E, Canner with rack. F, Tongs to fit bottles. G, Thermometer reading to 230°F (110°C). H, Fermentation lock.

harsh flavor because of too much alcohol. Two parts of sugar will usually produce about one part of alcohol and one part of carbon dioxide. A sugar content of 20% will therefore produce 11-12% alcohol in a completely fermented wine, the best concentration for table use. For dessert wines, which are sweeter, the alcohol may be increased to 14-15% by using a total sugar content of 24%.

Use a hydrometer to measure the sugar. The instrument should be calibrated in percent sugar (Balling or Brix scale) and

should cover a range from 0 to 35%. Hydrometers are available from stores that specialize in home wine makers' supplies or from scientific supply houses.

A simple calculation for the amount of sugar to add is this. Draw a rectangle. At the top left corner write the percent sugar content you have as measured by a hydrometer. At the bottom left write 100 (sugar is 100%). In the center write 20, the desired sugar percentage (24 for high-alcohol wines). Subtract the smaller value at top left from the center value and write the difference at bottom right. Subtract the center value from the larger value at bottom left and write the result at top right. The top right value is the proportion of juice and the bottom right



Figure 7. Small juice-extracting units. A and B, Juicers suitable for extracting several gallons of juice for canning or freezing. C. Cone-shaped food press.

value is the proportion of sugar to use to give the desired concentration. For example, if these values come out to 80 and 4 then for each 80 lb (36 kg) of juice or pulp add 4 lb (1.8 kg) of sugar or one part sugar to 20 parts of juice.

If the juice or pulp has been weighed accurately and the correct amount of sugar calculated to give 20 or 24%, the yeasts will convert all of this sugar to alcohol and carbon dioxide to give a dry wine.

When the production of gas slows down and finally stops, sediment begins to settle on the bottom of the carboy. Unlike other fruits, grapes contain a high proportion of tartaric acid, which gives the harsh flavor to new wine. The sediment consists of yeast cells and tartrates or wine stone.

Shortly after the fermenting stops, rack the wine by siphoning it carefully off the sediment into a clean carboy or glass jug. Always keep containers filled to the neck. To speed up the removal of tartrates, the wine may be chilled almost to the freezing point after fermentation is complete. Repeat the racking procedure at intervals until no more sediment forms. Racking also incorporates a small amount of air, which helps to hasten the aging process.

If the wine is too acid, a small amount of precipitated chalk may improve it. Add not more than $\frac{1}{4}$ teaspoon/gal ($1 \text{ cm}^3/4$ litres) a little at a time until the desired acidity is reached. Additional sediment will settle out if this is done.

Usually the wine will clear very well with these treatments, if it has been made properly. If cloudiness persists, fining with bentonite may clear the wine. Mix 1 teaspoon of the powder in a little water and stir into each gallon of wine ($1 \text{ cm}^3/\text{litre}$). Carefully remove the settled sediment by racking after 4 or 5 days. Bentonite is available at wine supply shops.

Gelatin has been used for fining, but it is not recommended because problems may arise from its use.

The wine should be brilliantly clear after these treatments and it is then ready for bottling. If it is too dry for the individual taste, sweeten it by adding sufficient sugar. One to 2% sugar will reduce the sour taste of an acid wine and improve its body.

To avoid the possibility of further fermentation in the bottle add 1 or 2 Campden tablets to each gallon (4.5 litres) before bottling. One-quarter teaspoon of ascorbic acid to each 2 gal ($1 \text{ cm}^3/8$ litres) will ensure that the color and flavor are maintained.

Fill the bottles to the neck and seal with new tight corks or new screw caps. Champagne-type bottles with plastic stoppers are very good. Bottles must be well sealed to prevent air from spoiling the wine. Thirteen-ounce (369-ml) or 26-ounce (738-ml) seal better than larger bottles.

Store bottles with cork stoppers on their sides to prevent the corks from drying out. Keep the wine in a cool dark place.

Aging

Wine may be stored in wooden barrels to age, but this is not essential if chilling and fining have been properly carried out after the fermentation. If you use a barrel, make sure it is clean and has no musty or vinegary odor. Barrels that have been used for brandy, whiskey, rum, or wine may be used for aging.

Always fill the barrel completely and do not leave the wine too long in the barrel. For barrels with a capacity of 10 gal (45 litres), store the wine for no longer than 6 weeks before bottling.

Wine matures in bottles, but more slowly than in a barrel. If a heavy deposit forms on standing, transfer the wine into clean bottles by pouring carefully to avoid disturbing the sediment. This treatment adds small amounts of air, which help to smooth a harsh-flavored wine. Do not partly fill bottles because this causes loss of color and sometimes spoilage, but fill them to the neck and store in a cool dark place.

Concentrated grape juices may be used where fresh fruit is not available. These may be purchased from stores specializing in wine-making materials. Directions for their use are given on the tins.

If you cannot make the wine soon after the grapes have been picked, you may store them. Stem and crush the grapes, add ascorbic acid and Campden tablets, and place them in the freezer until you are ready to proceed. Pressed juice for white wines may be treated in the same way.

FRUIT WINES

The general procedures for preparing fruit wines are similar to those for making grape wine. Because some fruits are high in flavor and acid, dilute the pulp or juice with water before fermenting.

Most fruits contain less natural yeast nutrients than grapes, and because the fruit juice may be further diluted with water, it is best to add some yeast nutrients. They are available from wine equipment suppliers, and usually come in powdered form

with directions for their use. If they are not available, 1 teaspoon of malt extract added to each gallon of juice or pulp (1 cm³/litre) is a good source of yeast nutrients.

The following procedures have produced acceptable wines. The suggested proportions of fruit to water have proved satisfactory, but the proportions may be varied to suit individual taste. Always use well-ripened fruit, but be sure it has no rot or mold.

Apricot, Peach, and Prune Wines

1. Halve, pit, and thoroughly mash the fruit.
2. Weigh the pulp, add an equal amount of water, and while stirring constantly, bring the mixture to a boil.
3. To each 6 lb (3 kg) of the mixture add 1 lb (0.5 kg) of sugar and cool to 100°F (38°C). Add 2 Campden tablets and ¼ teaspoon (1.2 cm³) of ascorbic acid to each gallon (4.5 litres) or 10 lb (4.5 kg) of pulp together with pectinase according to the suppliers directions. Mix well and allow to stand for 24 hr. Add yeast nutrient as directed on the package.
4. Inoculate with yeast as described for grape wine in the section "Yeasts and Fermentation."
5. After 48 hr, siphon the free-run juice from the pulp into a clean carboy, place the fermentation lock in position and ferment to completion.
6. Rack the wine and treat it in the same way as grape wine, except that no chilling is necessary.

Cherry Wine

1. Mash whole ripe cherries.
2. Proceed as directed from step 2 for apricot, peach, and prune wines.

Loganberry and Raspberry Wines

1. Mash ripe berries and add 1 gal of water for each 3 lb of

berries (5 litres/1.5 kg).

2. Add 1 lb (0.5 kg) of sugar for each pound (0.5 kg) of berries.
3. Add Campden tablets, ascorbic acid, and pectinase as in step 3 for apricot, peach, and prune wines.
4. Inoculate with yeast as described in the section "Yeasts and Fermentation" for grape wine.
5. Ferment for 7 days, siphon off the juice, and let it continue to ferment until finished.

Because the wine from both these fruits is high in flavor and acidity, it is most pleasant when slightly sweet. Add sugar before bottling, remembering to add also 2 Campden tablets and $\frac{1}{4}$ teaspoon of ascorbic acid to each gallon (1 cm³/4 litres).

Black Currant Wine

1. Remove the fruit from the stems and crush the berries.
2. Add $\frac{1}{2}$ gal of water for each pound of fruit (5 litres/kg).
3. Add 1 lb of sugar for each $\frac{1}{2}$ gal of this mixture (1 kg/5 litres).
4. Add 1 Campden tablet and pectinase to each $\frac{1}{2}$ gal (2.3 litres) of pulp, mix well, and let the mixture stand for 24 hr.
5. Proceed as for loganberry and raspberry wines. Ascorbic acid is not necessary because currants are high in this substance.

Dandelion Wine

1. To each 3 qt (3.4 litres) of dandelion flowers add 1 gal (4.5 litres) of water, and boil for 15 to 20 min.
2. Strain off the flowers, and to each gallon (4.5 litres) of liquid add 3 lb (1.4 kg) of sugar, 3 sliced lemons, 1 sliced orange, and yeast nutrient.
3. To each gallon (4.5 litres) of the resulting mixture add 2 Campden tablets, stir well, and let the mixture stand for 24 hr.

- Inoculate with wine yeast and ferment for 8 to 10 days. Siphon the clear liquid from the sediment. Continue fermentation, and rack at intervals until the wine is clear before bottling.

Parsnip Wine

- Wash the parsnips and cut them into small pieces.
- To each 4 lb of parsnips add 1 gal of water (2 kg add 5 litres), bring the mixture to a boil, and simmer for 20 to 30 min.
- To each gallon (4.5 litres) of pulp add 3 lb (1.4 kg) of sugar, 3 sliced oranges, 3 sliced lemons, and yeast nutrient.
- To each gallon (4.5 litres) of pulp add 2 Campden tablets, stir, and let the mixture stand for 24 hr.
- Inoculate as described for grape wine in the section "Yeasts and Fermentation."
- Siphon the clear juice from the pulp after 48 hr and allow it to ferment until finished.
- Rack the wine at intervals until it is clear, and then bottle it.

This recipe may be used for other vegetables such as beets, turnips, marrows, and carrots.

Rhubarb Wine

- To 10 lb (4.5 kg) of rhubarb cut in small pieces, add 2 gal (9 litres) of water and cook until soft (as stewed rhubarb).
- Strain off the pulp through cheesecloth.
- Add precipitated chalk to the juice until the acidity has been removed, as indicated by a very bland flavor. This removes undesirable oxalic acid. Allow the treated juice to stand until the sediment settles and siphon off the clear juice.
- Add Campden tables, ascorbic acid, and pectinase as for other wines.
- To each gallon add 4 lb of sugar (5 litres add 2 kg) and the

required amount of yeast nutrient.

6. Inoculate and ferment as described for grape wine in the section "Yeasts and Fermentation."
7. Before bottling, add citric acid to the wine to replace the oxalic acid removed by the chalk. Dissolve crystalline citric acid (obtained from the wine supply store) in a little water and add until the desired tartness is obtained.

If a sweet wine is desired, add more sugar and remember to add more Campden tablets to prevent bottle fermentation.

cider

Cider of the best quality is made from a blend of juices of different apple varieties. Varieties such as McIntosh or Winesap when used alone may give slightly unpleasant flavors. Jonathan yields a cider too high in acid for some tastes. Blends that have proved satisfactory are those containing 75% Delicious and 25% McIntosh or equal parts of Delicious and McIntosh, Jonathan, Newtown, or Winesap.

The following process yields an almost dry, lightly carbonated cider with an alcohol content slightly higher than that of beer.

Containers for Fermentation

Glass or plastic carboys or stoneware crocks are suitable containers in which to ferment the juice. During fermentation stopper carboys with a fermentation lock. Cover stoneware crocks with a suitable cloth such as fine cheesecloth.

Preparing the Juice

Strain the freshly pressed juice through cheesecloth to remove coarse particles of fruit. To each gallon (4.5 litres) of juice add 2 Campden tablets and pectinase as directed on the package, stir well, and let the mixture stand for 24 hr. This treatment prevents darkening of the cider and destroys wild yeasts and vinegar-producing bacteria.

Fermentation

Inoculate the juice with a wine yeast as described for grape wine in the section "Yeasts and Fermentation." The fermentation rate depends on the temperature and on the amount of sugar in the juice; fermentation should take 5 to 10 days. Nearly all the sugar will then have been used by the yeast. Completion of the fermentation may be judged by the slowing down of gas bubbles rising to the surface and the absence of sweetness. Leave the cider for 2 or 3 days to settle and clear, and then siphon it free from the sediment.

Before bottling, add 1 oz to $\frac{1}{2}$ lb (28 to 227 g) of sugar to each gallon (4.5 litres); $\frac{1}{2}$ lb gives a rather sweet cider. Fill the bottles leaving 1 to $1\frac{1}{2}$ in. (2.5–3.8 cm) of headspace, cap them, and hold them at 70°F (21°C) to allow bottle fermentation to take place. The carbon dioxide produced gives the cider sparkle or head. Open a trial bottle each day, and if enough carbonation has developed, the cider is ready to be pasteurized.

Pasteurization

Place the bottles upright on the bottom rack of a canner or boiler, and cover the bottles completely with water. Fill one uncapped bottle with water and raise it from the bottom so that the neck is above the water level. Put a thermometer into a cork so that the bulb is at the center of the bottle when the cork is in place resting on the neck of the bottle. Heat the water until the thermometer in the test bottle registers a temperature of 150°F (66°C); pasteurization is then complete.

Remove the bottles from the water and place them on their sides on several layers of paper. After 2 or 3 min place the bottles in warm water and leave them at room temperature until the cider is cool.

Storage

Store the containers of finished cider in an upright position in a cool place. The sediment should settle completely after 10 to 12 days, leaving clear liquid on the top. The cider should be ready to drink after about a month. When pouring the cider be careful

not to pour out any sediment, if you wish a clear sparkling glass.

Still Sweet Cider

To make a clear still cider add about $\frac{1}{2}$ lb of sugar to each gallon of freshly pressed juice (0.25 kg / 5 litres). Ferment until the degree of sweetness is to your taste and then stop the fermentation by adding 3 Campden tablets to each gallon (4.5 litres) and allow the cider to clear by settling and frequent racking in a cool place. Bottle when it is clear. Pasteurization is not necessary.

THINGS TO REMEMBER

1. Use a pure culture wine yeast.
2. Use Campden tablets.
3. Ferment and store wine in the dark.
4. Ferment at 70° F (21° C). Store at 50–60° F (10–16° C).
5. Do not leave wine in small barrels for more than 4-6 weeks. Keep full.
6. Use new tight corks, screw caps, or plastic stoppers.
7. Bottle in 13- or 26-oz (369- or 738-ml) bottles.

GLOSSARY OF TERMS

<i>Acetic</i>	<i>vinegar flavor</i>
<i>Brix degrees</i>	<i>percent sugar</i>
<i>Balling degrees</i>	<i>percent sugar</i>
<i>Dry</i>	<i>no sugar left</i>
<i>Lees</i>	<i>sediment</i>
<i>Must</i>	<i>crushed fruit or juice before or during fermentation</i>
<i>Oxidized</i>	<i>spoiled by exposure to air</i>
<i>Racking</i>	<i>separation of wine from pulp or lees</i>
<i>Semi sweet wine</i>	<i>with about 2–3% sugar</i>
<i>Sweet wine</i>	<i>with 6% sugar or more</i>

CONVERSION FACTORS FOR METRIC SYSTEM

Imperial units	Approximate conversion factor	Results in:
LINEAR		
inch	x 25	millimetre (mm)
foot	x 30	centimetre (cm)
yard	x 0.9	metre (m)
mile	x 1.6	kilometre (km)
AREA		
square inch	x 6.5	square centimetre (cm ²)
square foot	x 0.09	square metre (m ²)
acre	x 0.40	hectare (ha)
VOLUME		
cubic inch	x 16	cubic centimetre (cm ³)
cubic foot	x 28	cubic decimetre (dm ³)
cubic yard	x 0.8	cubic metre (m ³)
fluid ounce	x 28	millilitre (mℓ)
pint	x 0.57	litre (ℓ)
quart	x 1.1	litre (ℓ)
gallon	x 4.5	litre (ℓ)
bushel	x 0.36	hectolitre (hℓ)
WEIGHT		
ounce	x 28	gram (g)
pound	x 0.45	kilogram (kg)
short ton (2000 lb)	x 0.9	tonne (t)
TEMPERATURE		
degree fahrenheit	°F-32 x 0.56 (or °F-32 x 5/9)	degree Celsius (°C)
PRESSURE		
pounds per square inch	x 6.9	kilopascal (kPa)
POWER		
horsepower	x 746 x 0.75	watt (W) kilowatt (kW)
SPEED		
feet per second	x 0.30	metres per second (m/s)
miles per hour	x 1.6	kilometres per hour (km/h)
AGRICULTURE		
gallons per acre	x 11.23	litres per hectare (ℓ/ha)
quarts per acre	x 2.8	litres per hectare (ℓ/ha)
pints per acre	x 1.4	litres per hectare (ℓ/ha)
fluid ounces per acre	x 70	millilitres per hectare (mℓ/ha)
tons per acre	x 2.24	tonnes per hectare (t/ha)
pounds per acre	x 1.12	kilograms per hectare (kg/ha)
ounces per acre	x 70	grams per hectare (g/ha)
plants per acre	x 2.47	plants per hectare (plants/ha)

CAL/BCA OTTAWA K1A 0C5



3 9073 00234982 9

